

Some aspects of calculus on non-smooth sets

Stephen Semmes
Rice University

Let E be a closed set in \mathbf{R}^n , and suppose that there is a $k \geq 1$ such that every $x, y \in E$ can be connected by a rectifiable path in E with length $\leq k|x-y|$. This condition is satisfied by chord-arc curves, Lipschitz manifolds of any dimension, and fractals like Sierpinski gaskets and carpets. Note that length-minimizing paths in E are chord-arc curves with constant k .

A basic feature of this condition is that one can integrate local Lipschitz conditions on E to get global conditions. For instance, if $f : E \rightarrow \mathbf{R}$ is locally Lipschitz of order 1 with constant $C \geq 0$, then f is globally Lipschitz on E with constant kC .

Let $A(x)$ be a continuous function on E with values in linear mappings from \mathbf{R}^n to \mathbf{R} . Equivalently, one can use the standard inner product on \mathbf{R}^n to represent $A(x)$ by a continuous mapping from E into \mathbf{R}^n . Also let f be a locally Lipschitz real-valued function on E .

Suppose that A includes the directional derivatives of f almost everywhere on any rectifiable curve in E , in the sense that the derivative of $f(p(t))$ is equal to $A(p(t))$ applied to the derivative of $p(t)$ for almost every t when $p(t)$ is a locally Lipschitz function on an interval I in the real line with values in E . In particular, this holds when f is continuously differentiable on E in the sense of the Whitney extension theorem with differential at $x \in E$ given by $A(x)$. In this case,

$$(1) \quad f(p(b)) - f(p(a)) = \int_a^b A(p(t))(p'(t)) dt$$

for every $a, b \in I$. If $A(x)$ is constant on E , then it follows that f is equal to the restriction of an affine function on \mathbf{R}^n to E .

If A is not constant, then we still have that

$$(2) \quad |f(y) - f(x) - A(x)(y-x)| \leq k|x-y| \sup\{|A(w) - A(x)| : w \in E, |x-w| \leq k|x-y|\}$$

for every $x, y \in E$. This follows from the previous formula applied to a curve in E that connects x to y and has length $\leq k|x-y|$, by approximating $A(p(t))$ in the integral by $A(x)$ and estimating the remainder by the oscillation of A on the path times the length of the path. If we did not know already that f is continuously differentiable on E in Whitney's sense, then it follows from this and the continuity of A . Even if we did know that f is continuously-differentiable

on E , this estimate provides more precise information about the behavior of f in terms of the continuity properties of A . For example, Hölder continuity of order α of A implies $C^{1,\alpha}$ -type smoothness of f .

If E is a Lipschitz submanifold of \mathbf{R}^n , then E has tangent planes almost everywhere and a locally Lipschitz function f on E has linear differentials on almost all of these tangent planes. If there is a continuous function $A(x)$ on E with values in linear functionals on \mathbf{R}^n such that these differentials of f are equal to the restriction of $A(x)$ to the tangent plane to E at almost everywhere $x \in E$, then similar reasoning shows that f is continuously differentiable on E in Whitney's sense.

If E happens to be a C^1 submanifold of \mathbf{R}^n , then f is a C^1 function on E in the usual sense. Otherwise, restrictions of smooth functions on \mathbf{R}^n to E may not be smooth in a more intrinsic way. It may be that E is a non-smooth embedding of a smooth curve or surface, and that smooth functions on \mathbf{R}^n do not correspond to smooth functions on the parameter domain. It may be that E is something like a Lipschitz manifold for which there is not a compatible smooth structure. Nonetheless, we get some regularity to extrinsic smoothness under the geometric condition under consideration.

If E is a chord-arc curve, then one can integrate a continuous family $A(x)$ of differentials on E to get a function f . This is analogous to integrating a continuous function on an interval to get a C^1 function, but it is not quite the same, since the smoothness is now defined extrinsically.

Let us emphasize that this type of geometric condition does not imply anything like Poincaré or Sobolev inequalities, aside from the special case of chord-arc curves. For instance, one can have bubbles with small boundaries where a curve can easily pass. Thus one works with quantities based on the supremum norm instead of integrals.

Just as classical Fourier analysis on the real line deals with the ordinary differential operator d/dx , one can look at analysis on Lipschitz graphs in the complex plane as dealing with certain perturbations of d/dx , as in [4, 5]. The complex derivative of a holomorphic function on a neighborhood of the graph at a point on the graph is a multiple of the ordinary derivative tangent to the graph, which corresponds to a perturbation of d/dx when the graph is parameterized by projecting to the real line. This also works for a complex-valued continuously-differentiable function on the graph in Whitney's sense for which the differentials are complex-linear.

Additional conditions for the differentials of continuously-differentiable functions in Whitney's sense like complex-linearity can be very interesting. In particular, they may imply uniqueness of the differentials. As another version of this, a real-linear mapping from \mathbf{R}^n into a Clifford algebra is uniquely determined by its restriction to any hyperplane when it is left or right Clifford holomorphic. Thus it is natural to look at Clifford-valued continuously-differentiable functions on hypersurfaces in \mathbf{R}^n for which the differentials are left or right Clifford holomorphic. Note that the differentials of a continuously-differentiable function in Whitney's sense are uniquely determined by the function at any point where the set is not approximately contained in a hyperplane.

Normally, the differentials of a continuously-differentiable function in Whitney's sense are not uniquely determined by the function on the set, and they may not be very stable even if they are uniquely determined. The set may be contained in a smooth submanifold of lower dimension, so that the behavior of the differentials in the normal directions is not important. If there is suitable uniqueness and stability, then continuity properties of the differentials can be derived from the corresponding approximation properties of the function by affine functions on the set.

Uniqueness of the differential, perhaps through an additional condition like complex-linearity, has the effect of allowing a derivative in any direction to be interpreted as being tangent to the set. In this way, extrinsic behavior becomes more intrinsic.

References

- [1] L. Ambrosio and P. Tilli, *Topics on Analysis in Metric Spaces*, Oxford University Press, 2004.
- [2] F. Brackx, R. Delanghe, and F. Sommen, *Clifford Analysis*, Pitman, 1982.
- [3] J. Cheeger, *Differentiability of Lipschitz functions on metric measure spaces*, Geometric and Functional Analysis **9** (1999), 428–517.
- [4] R. Coifman and Y. Meyer, *Fourier analysis of multilinear convolutions, Calderón's theorem, and analysis of Lipschitz curves*, in *Euclidean Harmonic Analysis*, 104–122, Lecture Notes in Mathematics **779**, Springer-Verlag, 1980.
- [5] R. Coifman and Y. Meyer, *Nonlinear harmonic analysis, operator theory and P.D.E.*, in *Beijing Lectures in Harmonic Analysis*, Princeton University Press, 1986.
- [6] R. Coifman and G. Weiss, *Analyse Harmonique Non-Commutative sur Certains Espaces Homogènes*, Lecture Notes in Mathematics **242**, Springer-Verlag, 1971.
- [7] R. Coifman and G. Weiss, *Extensions of Hardy spaces and their use in analysis*, Bulletin of the American Mathematical Society **83** (1977), 569–645.
- [8] L. Evans and R. Gariepy, *Measure Theory and Fine Properties of Functions*, CRC Press, 1992.
- [9] K. Falconer, *The Geometry of Fractal Sets*, Cambridge University Press, 1986.
- [10] H. Federer, *Geometric Measure Theory*, Springer-Verlag, 1969.

- [11] B. Franchi, P. Hajłasz, and P. Koskela, *Definitions of Sobolev classes on metric spaces*, Annales de l'Institut Fourier (Grenoble) **49** (1999), 1903–1924.
- [12] M. Freedman and F. Quinn, *Topology of 4-Manifolds*, Princeton University Press, 1990.
- [13] J. Gilbert and M. Murray, *Clifford Algebras and Dirac Operators in Harmonic Analysis*, Cambridge University Press, 1991.
- [14] P. Hajłasz, *Sobolev spaces on an arbitrary metric space*, Potential Analysis **5** (1996), 403–415.
- [15] P. Hajłasz and P. Koskela, *Sobolev meets Poincaré*, Comptes Rendus de l'Académie des Sciences Série I Mathématique **320** (1995), 1211–1215.
- [16] P. Hajłasz and P. Koskela, *Sobolev Met Poincaré*, Memoirs of the American Mathematical Society **688**, 2000.
- [17] J. Heinonen, *Lectures on Analysis on Metric Spaces*, Springer-Verlag, 2001.
- [18] J. Heinonen, *Nonsmooth calculus*, Bulletin of the American Mathematical Society (N.S.) **44** (2007), 163–232.
- [19] J. Heinonen and P. Koskela, *Definitions of quasiconformality*, Inventiones Mathematicae **120** (1995), 61–79.
- [20] J. Heinonen and P. Koskela, *From local to global in quasiconformal structures*, Proceedings of the National Academy of Sciences USA **93** (1996), 554–556.
- [21] J. Heinonen and P. Koskela, *Quasiconformal maps in metric spaces with controlled geometry*, Acta Mathematica **181** (1998), 1–61.
- [22] J. Heinonen and P. Koskela, *A note on Lipschitz functions, upper gradients, and the Poincaré inequality*, New Zealand Journal of Mathematics **28** (1999), 37–42.
- [23] J. Heinonen and D. Sullivan, *On the locally branched Euclidean metric guage*, Duke Mathematical Journal **114** (2002), 15–41.
- [24] J. Kigami, *Analysis on Fractals*, Cambridge University Press, 2001.
- [25] R. Kirby and L. Siebenmann, *Foundational Essays on Topological Manifolds, Smoothings, and Triangulations*, Princeton University Press, 1977.
- [26] S. Krantz, *A Panorama of Harmonic Analysis*, Mathematical Association of America, 1999.
- [27] S. Krantz and H. Parks, *The Geometry of Domains in Space*, Birkhäuser, 1999.

- [28] P. Mattila, *Geometry of Sets and Measures in Euclidean Spaces*, Cambridge University Press, 1995.
- [29] J. Luukkainen and J. Väisälä, *Elements of Lipschitz Topology*, Annales Academiae Scientiarum Fennicae Series A I Mathematica **3** (1977), 85–122.
- [30] A. McIntosh, *Clifford algebras, Fourier theory, singular integrals, and harmonic functions on Lipschitz domains*, in *Clifford Analysis and Related Topics*, 33–87, CRC Press, 1996.
- [31] F. Morgan, *Geometric Measure Theory: A Beginner's Guide*, 3rd edition, Academic Press, 2000.
- [32] S. Rickman, *Quasiregular Mappings*, Springer-Verlag, 1993.
- [33] N. Shanmugalingam, *Newtonian spaces: An extension of Sobolev spaces to metric measure spaces*, Revista Matemática Iberoamericana **16** (2000), 243–279.
- [34] E. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, 1970.
- [35] E. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, with the assistance of T. Murphy, Princeton University Press, 1993.
- [36] E. Stein and G. Weiss, *Introduction to Fourier Analysis on Euclidean Spaces*, Princeton University Press, 1971.
- [37] R. Strichartz, *Differential Equations on Fractals*, Princeton University Press, 2006.
- [38] D. Sullivan, *Hyperbolic geometry and homeomorphisms*, in *Geometric Topology*, 543–555, Academic Press, 1979.
- [39] D. Sullivan, *Exterior d , the local degree, and smoothability*, in *Prospects in topology*, 328–338, Princeton University Press, 1995.
- [40] H. Whitney, *Geometric Integration Theory*, Princeton University Press, 1957.